

## List of Figures

1	Decay of $\text{Ti}(\text{NRAr}^1)_3$ ( <b>1</b> ) monitored by UV-vis spectroscopy: reaction with a 1,4-diketone (top); reaction with bromo-benzene – typical decay (bottom). . . . .	3
2	EPR spectrum of <b>1</b> -OCPh <sub>2</sub> , toluene, 25 °C). . . . .	4
3	<sup>1</sup> H NMR spectra (CDCl <sub>3</sub> ) of the Gomberg-like dimer ( <b>1</b> -OCMePh) <sub>2</sub> (top) and its disproportionation products, enolate <b>1</b> -OC(CH <sub>2</sub> )Ph and alkoxide <b>1</b> -OCHMePh (bottom). . . . .	5
4	Infrared spectra (C <sub>6</sub> D <sub>6</sub> , KBr plates) of succinic acid di-amide (top), amide enolate <b>1</b> -OC(CH <sub>2</sub> )NPhMe (center) and aldol condensation product <b>1</b> -OCHPh(CH <sub>2</sub> CONPhMe) (bottom). . . . .	6
5	Frontier molecular orbitals of <b>2</b> -OCHCH <sub>2</sub> (top) and <b>2</b> -OC(NH <sub>2</sub> )CH <sub>2</sub> (bottom). . . . .	24

## List of Tables

1	Atomic coordinates [ $\times 10^4$ ] and equivalent isotropic displacement parameters [ $\text{\AA}^2 \times 10^3$ ] for <b>1</b> -OC(CH <sub>2</sub> )NPhMe. $U(\text{eq})$ is defined as one third of the trace of the orthogonalized $U_{ij}$ tensor. . . . .	7
2	Bond lengths [ $\text{\AA}$ ] and angles [ $^\circ$ ] for <b>1</b> -OC(CH <sub>2</sub> )NPhMe. . . . .	9
3	Anisotropic displacement parameters [ $\text{\AA}^2 \times 10^3$ ] for <b>1</b> -OC(CH <sub>2</sub> )NPhMe. . . . .	13
4	Atomic coordinates [ $\times 10^4$ ] and equivalent isotropic displacement parameters [ $\text{\AA}^2 \times 10^3$ ] for <b>1</b> -OC(CH <sub>2</sub> )Ar <sup>2</sup> . $U(\text{eq})$ is defined as one third of the trace of the orthogonalized $U_{ij}$ tensor. . . . .	15
5	Bond lengths [ $\text{\AA}$ ] and angles [ $^\circ$ ] for <b>1</b> -OC(CH <sub>2</sub> )Ar <sup>2</sup> . . . . .	17
6	Anisotropic displacement parameters [ $\text{\AA}^2 \times 10^3$ ] for <b>1</b> -OC(CH <sub>2</sub> )Ar <sup>2</sup> . . . . .	21
7	Structural parameters for <b>1</b> -OC(CH <sub>2</sub> )Ar <sup>2</sup> and <b>1</b> -OC(CH <sub>2</sub> )NPhMe (crystallographically determined) and <b>2</b> -OCHCH <sub>2</sub> and <b>2</b> -OC(NH <sub>2</sub> )CH <sub>2</sub> (calculated). . . . .	23

## Contents

1	DFT calculation for <b>2</b> -OC(NH <sub>2</sub> )CH <sub>2</sub>	25
---	---	----

Agapie et al.

*Radical Scission of Symmetrical 1,4-Dicarbonyl Compounds*

3

**2 DFT calculation for 2-OCHCH<sub>2</sub>**

29




Figure 1: Decay of  $\text{Ti}(\text{NRAr}^1)_3$  (1) monitored by UV-vis spectroscopy: reaction with a 1,4-diketone (top); reaction with bromo-benzene – typical decay (bottom).




Figure 2: EPR spectrum of 1-OCPh<sub>2</sub>, toluene, 25 °C).




Figure 3:  $^1\text{H}$  NMR spectra ( $\text{CDCl}_3$ ) of the Gomberg-like dimer  $(1\text{-OCMePh})_2$  (top) and its disproportionation products, enolate  $1\text{-OC}(\text{CH}_2)\text{Ph}$  and alkoxide  $1\text{-OCHMePh}$  (bottom).




Figure 4: Infrared spectra ( $\text{C}_6\text{D}_6$ , KBr plates) of succinic acid di-amide (top), amide enolate **16** ( $\text{OC}(\text{CH}_2)\text{NPhMe}$ ) (center) and aldol condensation product **17** ( $\text{OCHPh}(\text{CH}_2\text{CONPhMe})$ ) (bottom).

Table 1: Atomic coordinates [ $\times 10^4$ ] and equivalent isotropic displacement parameters [ $\text{\AA}^2 \times 10^3$ ] for 1-OC(CH<sub>2</sub>)NPhMe.  $U(\text{eq})$  is defined as one third of the trace of the orthogonalized  $U_{ij}$  tensor.

	x	y	z	U(eq)
Ti	783(1)	1605(1)	2314(1)	27(1)
O	2014(2)	207(2)	2147(2)	35(1)
N(2)	-663(3)	1469(2)	3007(2)	30(1)
N(3)	1499(3)	2619(2)	2837(2)	29(1)
C(31)	1140(3)	3791(3)	2592(2)	31(1)
N(1)	177(3)	2259(3)	1258(2)	30(1)
C(310)	2460(4)	1208(3)	3914(3)	43(1)
C(11)	-1171(4)	2767(4)	1168(2)	35(1)
C(35)	1579(4)	5342(3)	1841(3)	41(1)
C(46)	3594(5)	-1928(4)	4183(3)	59(1)
C(39)	2298(5)	3201(4)	4111(3)	68(2)
C(21)	-1397(4)	2423(3)	3525(2)	33(1)
C(34)	407(4)	6057(3)	2153(3)	42(1)
C(44)	5496(5)	-1476(5)	4028(3)	69(2)
N(4)	4052(3)	-1097(3)	1990(2)	51(1)
C(42)	4272(4)	-1345(3)	2839(3)	41(1)
C(48)	5065(5)	-1021(5)	1415(3)	75(2)
C(27)	-1078(4)	440(3)	3155(2)	37(1)
C(411)	2415(5)	-1218(4)	1145(3)	51(1)
C(22)	-1027(4)	2580(3)	4315(2)	38(1)
C(19)	2395(4)	1776(4)	643(3)	44(1)
C(17)	997(4)	2280(3)	460(2)	36(1)
C(210)	-656(5)	-92(4)	3994(3)	57(1)
C(32)	-26(4)	4534(3)	2892(2)	38(1)

C(47)	3408(4)	-1709(4)	3341(3)	47(1)
C(37)	2521(4)	2268(3)	3457(2)	35(1)
C(33)	-407(4)	5673(3)	2676(3)	43(1)
C(36)	1931(4)	4210(3)	2065(2)	39(1)
C(29)	-2515(4)	746(4)	3140(3)	64(1)
C(43)	5326(4)	-1242(4)	3189(3)	52(1)
C(41)	2784(4)	-715(3)	1739(2)	37(1)
C(351)	2473(5)	5771(4)	1266(3)	65(2)
C(38)	3850(4)	2004(4)	3006(3)	60(1)
C(331)	-1675(4)	6468(4)	3031(4)	72(2)
C(24)	-2804(4)	4275(4)	4519(3)	51(1)
C(23)	-1715(4)	3503(4)	4817(3)	46(1)
C(28)	-463(4)	-421(4)	2461(3)	51(1)
C(18)	708(4)	1560(4)	-213(2)	47(1)
C(26)	-2496(4)	3210(3)	3253(3)	41(1)
C(110)	760(4)	3484(4)	135(3)	48(1)
C(25)	-3213(4)	4134(4)	3747(3)	48(1)
C(12)	-1766(4)	3932(4)	1231(2)	44(1)
C(16)	-1923(4)	2098(4)	1019(2)	43(1)
C(15)	-3227(4)	2558(5)	957(3)	55(1)
C(13)	-3063(4)	4426(4)	1128(3)	55(1)
C(14)	-3774(5)	3719(5)	1003(3)	65(2)
C(45)	4641(5)	-1813(5)	4528(3)	67(2)
C(231)	-1262(5)	3640(4)	5669(3)	70(2)
C(251)	-4426(5)	4959(4)	3449(3)	73(2)
C(131)	-3671(5)	5696(5)	1157(4)	84(2)
C(151)	-4045(5)	1821(5)	855(3)	79(2)



Table 2: Bond lengths [Å] and angles [°] for 1-OC(CH<sub>2</sub>)NPhMe.

Ti-O	1.847(3)
Ti-N(3)	1.931(3)
Ti-N(1)	1.931(3)
Ti-N(2)	1.936(3)
O-C(41)	1.340(4)
N(2)-C(21)	1.444(5)
N(2)-C(27)	1.510(5)
N(3)-C(31)	1.444(5)
N(3)-C(37)	1.510(5)
C(31)-C(32)	1.385(5)
C(31)-C(36)	1.390(5)
N(1)-C(11)	1.440(5)
N(1)-C(17)	1.519(4)
C(310)-C(37)	1.520(5)
C(11)-C(16)	1.392(5)
C(11)-C(12)	1.394(6)
C(35)-C(34)	1.383(6)
C(35)-C(36)	1.389(5)
C(35)-C(351)	1.520(6)
C(46)-C(45)	1.370(7)
C(46)-C(47)	1.386(6)
C(39)-C(37)	1.530(6)
C(21)-C(26)	1.388(5)
C(21)-C(22)	1.393(5)
C(34)-C(33)	1.380(6)
C(44)-C(45)	1.362(7)
C(44)-C(43)	1.381(6)
N(4)-C(42)	1.408(5)
N(4)-C(41)	1.417(5)
N(4)-C(48)	1.427(5)
C(42)-C(43)	1.377(6)
C(42)-C(47)	1.389(6)
C(27)-C(28)	1.523(5)
C(27)-C(29)	1.517(6)
C(27)-C(210)	1.523(6)
C(411)-C(41)	1.309(5)

---

C(22)-C(23)	1.391(5)
C(19)-C(17)	1.520(5)
C(17)-C(18)	1.531(5)
C(17)-C(110)	1.530(6)
C(32)-C(33)	1.393(5)
C(37)-C(38)	1.531(6)
C(33)-C(331)	1.513(6)
C(24)-C(25)	1.379(6)
C(24)-C(23)	1.389(6)
C(23)-C(231)	1.519(6)
C(26)-C(25)	1.393(6)
C(25)-C(251)	1.510(6)
C(12)-C(13)	1.390(6)
C(16)-C(15)	1.384(6)
C(15)-C(14)	1.381(7)
C(15)-C(151)	1.507(7)
C(13)-C(14)	1.387(7)
C(13)-C(131)	1.511(7)
O-Ti-N(3)	109.85(12)
O-Ti-N(1)	110.59(12)
N(3)-Ti-N(1)	107.84(13)
O-Ti-N(2)	110.51(12)
N(3)-Ti-N(2)	109.90(12)
N(1)-Ti-N(2)	108.08(12)
C(41)-O-Ti	159.0(2)
C(21)-N(2)-C(27)	113.2(3)
C(21)-N(2)-Ti	117.9(2)
C(27)-N(2)-Ti	128.6(2)
C(31)-N(3)-C(37)	114.9(3)
C(31)-N(3)-Ti	120.0(2)
C(37)-N(3)-Ti	124.9(2)
C(32)-C(31)-C(36)	118.3(4)
C(32)-C(31)-N(3)	119.4(3)
C(36)-C(31)-N(3)	122.3(3)
C(11)-N(1)-C(17)	113.6(3)
C(11)-N(1)-Ti	120.2(2)
C(17)-N(1)-Ti	126.2(2)
C(16)-C(11)-C(12)	118.0(4)
C(16)-C(11)-N(1)	120.5(4)
C(12)-C(11)-N(1)	121.5(4)
C(34)-C(35)-C(36)	118.3(4)

---

---

C(34)-C(35)-C(351)	121.5(4)
C(36)-C(35)-C(351)	120.2(4)
C(45)-C(46)-C(47)	120.8(5)
C(26)-C(21)-C(22)	117.9(4)
C(26)-C(21)-N(2)	121.0(3)
C(22)-C(21)-N(2)	121.0(3)
C(35)-C(34)-C(33)	121.7(4)
C(45)-C(44)-C(43)	121.7(5)
C(42)-N(4)-C(41)	119.6(3)
C(42)-N(4)-C(48)	120.2(4)
C(41)-N(4)-C(48)	119.1(4)
C(43)-C(42)-N(4)	121.8(4)
C(43)-C(42)-C(47)	118.6(4)
N(4)-C(42)-C(47)	119.6(4)
N(2)-C(27)-C(28)	109.2(3)
N(2)-C(27)-C(29)	111.0(3)
C(28)-C(27)-C(29)	107.9(3)
N(2)-C(27)-C(210)	110.2(3)
C(28)-C(27)-C(210)	108.9(4)
C(29)-C(27)-C(210)	109.6(4)
C(23)-C(22)-C(21)	121.6(4)
C(19)-C(17)-N(1)	108.7(3)
C(19)-C(17)-C(18)	108.8(3)
N(1)-C(17)-C(18)	109.4(3)
C(19)-C(17)-C(110)	108.7(3)
N(1)-C(17)-C(110)	111.7(3)
C(18)-C(17)-C(110)	109.4(3)
C(33)-C(32)-C(31)	121.3(4)
C(46)-C(47)-C(42)	120.2(4)
N(3)-C(37)-C(39)	110.0(3)
N(3)-C(37)-C(310)	109.8(3)
C(39)-C(37)-C(310)	107.8(3)
N(3)-C(37)-C(38)	110.7(3)
C(39)-C(37)-C(38)	110.5(4)
C(310)-C(37)-C(38)	108.1(3)
C(34)-C(33)-C(32)	118.7(4)
C(34)-C(33)-C(331)	121.3(4)
C(32)-C(33)-C(331)	120.0(4)
C(35)-C(36)-C(31)	121.7(4)
C(42)-C(43)-C(44)	120.0(4)
C(411)-C(41)-O	123.4(4)

---

C(411)-C(41)-N(4)	122.1(4)
O-C(41)-N(4)	114.5(3)
C(25)-C(24)-C(23)	121.3(4)
C(24)-C(23)-C(22)	118.6(4)
C(24)-C(23)-C(231)	121.9(4)
C(22)-C(23)-C(231)	119.5(4)
C(21)-C(26)-C(25)	121.6(4)
C(24)-C(25)-C(26)	118.9(4)
C(24)-C(25)-C(251)	120.5(4)
C(26)-C(25)-C(251)	120.6(4)
C(13)-C(12)-C(11)	121.6(4)
C(11)-C(16)-C(15)	121.8(4)
C(14)-C(15)-C(16)	118.3(4)
C(14)-C(15)-C(151)	120.3(5)
C(16)-C(15)-C(151)	121.4(5)
C(14)-C(13)-C(12)	118.0(5)
C(14)-C(13)-C(131)	121.5(5)
C(12)-C(13)-C(131)	120.5(5)
C(13)-C(14)-C(15)	122.1(4)
C(44)-C(45)-C(46)	118.7(5)

Table 3: Anisotropic displacement parameters [ $\text{\AA}^2 \times 10^3$ ] for 1-OC(CH<sub>2</sub>)NPhMe.

	U11	U22	U33	U23	U13	U12
Ti	27(1)	28(1)	27(1)	0(1)	-2(1)	-10(1)
O	37(2)	31(2)	35(2)	-2(1)	-3(1)	-8(1)
N(2)	33(2)	32(2)	25(2)	-1(1)	-3(1)	-12(2)
N(3)	27(2)	27(2)	31(2)	1(1)	-5(1)	-9(1)
C(31)	29(2)	34(2)	32(2)	2(2)	-10(2)	-11(2)
N(1)	26(2)	38(2)	26(2)	0(1)	-1(1)	-11(2)
C(310)	50(3)	38(3)	38(2)	1(2)	-8(2)	-8(2)
C(11)	29(2)	48(3)	27(2)	2(2)	-1(2)	-11(2)
C(35)	43(3)	36(3)	45(3)	5(2)	-6(2)	-14(2)
C(46)	55(3)	54(3)	59(3)	18(2)	5(3)	-7(3)
C(39)	96(4)	39(3)	68(3)	-4(2)	-47(3)	-9(3)
C(21)	33(2)	33(2)	33(2)	-1(2)	6(2)	-14(2)
C(34)	45(3)	26(2)	53(3)	2(2)	-11(2)	-8(2)
C(44)	42(3)	92(4)	64(4)	-5(3)	-11(3)	-8(3)
N(4)	36(2)	69(3)	43(2)	1(2)	7(2)	-10(2)
C(42)	31(2)	40(3)	45(3)	-3(2)	2(2)	-3(2)
C(48)	43(3)	112(5)	62(3)	10(3)	12(3)	-20(3)
C(27)	44(3)	37(2)	35(2)	-3(2)	2(2)	-23(2)
C(411)	61(3)	49(3)	45(3)	-16(2)	5(2)	-21(2)
C(22)	39(2)	37(2)	36(2)	-2(2)	1(2)	-11(2)
C(19)	35(2)	53(3)	41(2)	5(2)	2(2)	-15(2)
C(17)	30(2)	49(3)	27(2)	3(2)	1(2)	-13(2)
C(210)	86(4)	45(3)	48(3)	4(2)	-2(3)	-34(3)
C(32)	31(2)	38(3)	45(2)	2(2)	-2(2)	-11(2)
C(47)	39(3)	49(3)	51(3)	8(2)	-1(2)	-11(2)
C(37)	39(2)	30(2)	36(2)	-1(2)	-10(2)	-10(2)
C(33)	36(2)	32(2)	56(3)	-1(2)	-2(2)	-4(2)
C(36)	35(2)	36(2)	43(2)	1(2)	-1(2)	-9(2)
C(29)	48(3)	67(3)	88(4)	-9(3)	4(3)	-33(3)
C(43)	35(3)	63(3)	54(3)	-4(2)	2(2)	-10(2)
C(41)	35(2)	37(2)	39(2)	-2(2)	3(2)	-13(2)
C(351)	58(3)	47(3)	87(4)	22(3)	12(3)	-18(3)
C(38)	35(3)	75(4)	73(3)	25(3)	-20(2)	-19(2)
C(331)	51(3)	42(3)	109(5)	4(3)	11(3)	2(2)

C(24)	54(3)	40(3)	50(3)	-11(2)	19(2)	-8(2)
C(23)	58(3)	40(3)	39(2)	-10(2)	7(2)	-18(2)
C(28)	66(3)	43(3)	52(3)	-10(2)	7(2)	-30(2)
C(18)	49(3)	62(3)	31(2)	-6(2)	3(2)	-20(2)
C(26)	37(3)	45(3)	37(2)	2(2)	4(2)	-11(2)
C(110)	45(3)	56(3)	43(3)	15(2)	-2(2)	-18(2)
C(25)	35(3)	44(3)	56(3)	3(2)	14(2)	-4(2)
C(12)	41(3)	55(3)	34(2)	3(2)	0(2)	-15(2)
C(16)	36(3)	65(3)	30(2)	2(2)	0(2)	-22(2)
C(15)	37(3)	96(4)	35(3)	-3(3)	0(2)	-27(3)
C(13)	36(3)	69(4)	47(3)	14(2)	5(2)	0(3)
C(14)	29(3)	110(5)	46(3)	13(3)	-1(2)	-11(3)
C(45)	50(3)	89(4)	49(3)	12(3)	-8(3)	-4(3)
C(231)	92(4)	57(3)	52(3)	-23(3)	-1(3)	-13(3)
C(251)	48(3)	72(4)	76(4)	3(3)	12(3)	7(3)
C(131)	55(3)	84(4)	85(4)	23(3)	9(3)	13(3)
C(151)	54(3)	144(6)	58(3)	-10(3)	0(3)	-57(4)

Table 4: Atomic coordinates [ $\times 10^4$ ] and equivalent isotropic displacement parameters [ $\text{\AA}^2 \times 10^3$ ] for 1-OC(CH<sub>2</sub>)Ar<sup>2</sup>.  $U(\text{eq})$  is defined as one third of the trace of the orthogonalized  $U_{ij}$  tensor.

	x	y	z	U(eq)
O	5339(1)	3173(2)	1668(1)	35(1)
Ti(1)	6492(1)	2846(1)	1670(1)	27(1)
N(1)	6540(2)	2012(2)	897(1)	31(1)
C(1)	4695(3)	3274(3)	2614(2)	46(1)
N(2)	7129(2)	4065(2)	1640(1)	31(1)
C(2)	4624(2)	3181(3)	1962(2)	34(1)
N(3)	6999(2)	2151(2)	2471(1)	29(1)
C(3)	3764(2)	3091(3)	1495(2)	36(1)
C(4)	3614(2)	3531(3)	876(2)	46(1)
C(5)	2800(3)	3429(4)	453(2)	60(1)
C(6)	2134(3)	2880(4)	625(3)	62(1)
C(7)	2284(3)	2456(4)	1246(3)	68(1)
C(8)	3089(3)	2553(3)	1677(2)	57(1)
C(9)	1272(3)	2735(5)	140(3)	101(2)
C(11)	7185(2)	1243(3)	969(2)	33(1)
C(12)	6964(2)	239(3)	944(2)	39(1)
C(13)	7601(3)	-490(3)	1051(2)	43(1)
C(14)	8476(3)	-215(3)	1172(2)	46(1)
C(15)	8723(2)	769(3)	1192(2)	42(1)
C(16)	8068(2)	1486(3)	1088(2)	36(1)
C(17)	5839(2)	2003(3)	271(2)	40(1)
C(18)	5592(3)	3076(3)	88(2)	49(1)
C(19)	5016(2)	1449(3)	370(2)	53(1)
C(21)	7926(2)	4046(2)	1377(2)	33(1)
C(22)	7917(2)	4104(3)	698(2)	37(1)
C(23)	8685(3)	4106(3)	447(2)	45(1)
C(24)	9477(3)	4071(3)	891(2)	50(1)
C(25)	9518(2)	4014(3)	1572(2)	47(1)
C(26)	8739(2)	3991(3)	1811(2)	40(1)

C(27)	6834(3)	5100(3)	1777(2)	46(1)
C(28)	6349(3)	5085(3)	2345(3)	68(1)
C(29)	7583(4)	5808(4)	1928(4)	108(3)
C(31)	7786(2)	2492(2)	2903(2)	32(1)
C(32)	7287(2)	358(3)	2751(2)	42(1)
C(33)	8490(3)	3569(3)	3797(2)	48(1)
C(34)	9291(3)	3138(3)	3758(2)	50(1)
C(35)	9359(2)	2394(3)	3305(2)	45(1)
C(36)	8603(2)	2090(3)	2872(2)	39(1)
C(37)	6629(2)	1210(2)	2709(2)	32(1)
C(38)	5793(2)	935(3)	2213(2)	39(1)
C(39)	6396(3)	1368(3)	3392(2)	46(1)
C(110)	6191(3)	1551(4)	-313(2)	62(1)
C(131)	7326(3)	-1560(3)	1053(3)	71(1)
C(151)	9679(3)	1061(4)	1307(3)	68(1)
C(210)	6177(5)	5460(4)	1155(3)	114(3)
C(231)	8650(3)	4138(4)	-298(2)	68(1)
C(251)	10397(3)	3985(4)	2051(3)	75(2)
C(310)	7740(2)	3225(3)	3374(2)	40(1)
C(331)	8425(3)	4377(4)	4299(2)	70(1)
C(351)	10230(3)	1901(4)	3283(3)	68(1)



Table 5: Bond lengths [Å] and angles [°] for 1-OC(CH<sub>2</sub>)Ar<sup>2</sup>.

O-C(2)	1.352(4)
O-Ti(1)	1.826(2)
Ti(1)-N(3)	1.904(3)
Ti(1)-N(2)	1.920(3)
Ti(1)-N(1)	1.943(3)
N(1)-C(11)	1.422(4)
N(1)-C(17)	1.491(4)
C(1)-C(2)	1.311(5)
N(2)-C(21)	1.432(4)
N(2)-C(27)	1.511(4)
C(2)-C(3)	1.470(5)
N(3)-C(31)	1.422(4)
N(3)-C(37)	1.508(4)
C(3)-C(4)	1.366(5)
C(3)-C(8)	1.375(5)
C(4)-C(5)	1.375(6)
C(5)-C(6)	1.364(6)
C(6)-C(7)	1.361(7)
C(6)-C(9)	1.497(6)
C(7)-C(8)	1.372(6)
C(11)-C(16)	1.371(5)
C(11)-C(12)	1.394(5)
C(12)-C(13)	1.373(5)
C(13)-C(14)	1.370(5)
C(13)-C(131)	1.504(6)
C(14)-C(15)	1.379(6)
C(15)-C(16)	1.382(5)
C(15)-C(151)	1.493(5)
C(17)-C(19)	1.517(5)
C(17)-C(110)	1.524(5)
C(17)-C(18)	1.524(5)
C(21)-C(22)	1.376(5)
C(21)-C(26)	1.380(5)
C(22)-C(23)	1.376(5)
C(23)-C(24)	1.363(6)
C(23)-C(231)	1.502(6)
C(24)-C(25)	1.372(6)

C(25)-C(26)	1.379(5)
C(25)-C(251)	1.499(6)
C(27)-C(29)	1.481(6)
C(27)-C(28)	1.493(5)
C(27)-C(210)	1.529(7)
C(31)-C(36)	1.381(5)
C(31)-C(310)	1.386(5)
C(32)-C(37)	1.521(5)
C(33)-C(310)	1.373(5)
C(33)-C(34)	1.378(6)
C(33)-C(331)	1.509(6)
C(34)-C(35)	1.378(6)
C(35)-C(36)	1.373(5)
C(35)-C(351)	1.504(6)
C(37)-C(38)	1.512(5)
C(37)-C(39)	1.514(5)
C(2)-O-Ti(1)	151.2(2)
O-Ti(1)-N(3)	110.60(11)
O-Ti(1)-N(2)	107.12(11)
N(3)-Ti(1)-N(2)	108.88(12)
O-Ti(1)-N(1)	109.36(11)
N(3)-Ti(1)-N(1)	109.23(12)
N(2)-Ti(1)-N(1)	111.64(12)
C(11)-N(1)-C(17)	116.9(3)
C(11)-N(1)-Ti(1)	118.5(2)
C(17)-N(1)-Ti(1)	123.7(2)
C(21)-N(2)-C(27)	113.3(3)
C(21)-N(2)-Ti(1)	118.5(2)
C(27)-N(2)-Ti(1)	127.7(2)
C(1)-C(2)-O	122.3(3)
C(1)-C(2)-C(3)	122.8(3)
O-C(2)-C(3)	114.9(3)
C(31)-N(3)-C(37)	113.9(3)
C(31)-N(3)-Ti(1)	121.3(2)
C(37)-N(3)-Ti(1)	124.8(2)
C(4)-C(3)-C(8)	118.5(4)
C(4)-C(3)-C(2)	121.5(3)
C(8)-C(3)-C(2)	120.0(3)
C(3)-C(4)-C(5)	119.7(4)
C(6)-C(5)-C(4)	122.1(4)
C(7)-C(6)-C(5)	117.9(4)

---

C(7)-C(6)-C(9)	121.1(5)
C(5)-C(6)-C(9)	121.0(5)
C(6)-C(7)-C(8)	121.0(4)
C(7)-C(8)-C(3)	120.8(4)
C(16)-C(11)-C(12)	117.6(3)
C(16)-C(11)-N(1)	119.3(3)
C(12)-C(11)-N(1)	123.0(3)
C(13)-C(12)-C(11)	121.9(3)
C(14)-C(13)-C(12)	118.6(4)
C(14)-C(13)-C(131)	121.7(4)
C(12)-C(13)-C(131)	119.7(4)
C(13)-C(14)-C(15)	121.4(4)
C(14)-C(15)-C(16)	118.7(3)
C(14)-C(15)-C(151)	121.0(4)
C(16)-C(15)-C(151)	120.3(4)
C(11)-C(16)-C(15)	121.7(3)
N(1)-C(17)-C(19)	111.8(3)
N(1)-C(17)-C(110)	110.9(3)
C(19)-C(17)-C(110)	109.9(3)
N(1)-C(17)-C(18)	107.6(3)
C(19)-C(17)-C(18)	109.0(3)
C(110)-C(17)-C(18)	107.5(3)
C(22)-C(21)-C(26)	118.0(3)
C(22)-C(21)-N(2)	122.2(3)
C(26)-C(21)-N(2)	119.8(3)
C(23)-C(22)-C(21)	122.1(4)
C(24)-C(23)-C(22)	118.3(4)
C(24)-C(23)-C(231)	120.9(4)
C(22)-C(23)-C(231)	120.7(4)
C(23)-C(24)-C(25)	121.5(4)
C(24)-C(25)-C(26)	119.1(4)
C(24)-C(25)-C(251)	120.5(4)
C(26)-C(25)-C(251)	120.3(4)
C(25)-C(26)-C(21)	120.9(4)
C(29)-C(27)-C(28)	109.7(4)
C(29)-C(27)-N(2)	112.6(3)
C(28)-C(27)-N(2)	110.1(3)
C(29)-C(27)-C(210)	109.0(5)
C(28)-C(27)-C(210)	107.0(4)

---

---

N(2)-C(27)-C(210)	108.4(4)
C(36)-C(31)-C(310)	118.9(3)
C(36)-C(31)-N(3)	121.0(3)
C(310)-C(31)-N(3)	120.1(3)
C(310)-C(33)-C(34)	118.1(4)
C(310)-C(33)-C(331)	120.3(4)
C(34)-C(33)-C(331)	121.6(4)
C(35)-C(34)-C(33)	122.2(4)
C(36)-C(35)-C(34)	118.5(4)
C(36)-C(35)-C(351)	119.6(4)
C(34)-C(35)-C(351)	121.9(4)
C(35)-C(36)-C(31)	121.1(4)
N(3)-C(37)-C(38)	108.6(3)
N(3)-C(37)-C(39)	110.3(3)
C(38)-C(37)-C(39)	108.5(3)
N(3)-C(37)-C(32)	111.4(3)
C(38)-C(37)-C(32)	108.6(3)
C(39)-C(37)-C(32)	109.3(3)
C(33)-C(310)-C(31)	121.3(4)

---

Table 6: Anisotropic displacement parameters [ $\text{\AA}^2 \times 10^3$ ] for 1-OC(CH<sub>2</sub>)Ar<sup>2</sup>.

	U11	U22	U33	U23	U13	U12
O	31(1)	36(1)	38(1)	-1(1)	7(1)	0(1)
Ti(1)	27(1)	27(1)	27(1)	0(1)	5(1)	0(1)
N(1)	30(2)	31(2)	30(2)	-2(1)	3(1)	3(1)
C(1)	43(2)	51(2)	46(2)	3(2)	12(2)	7(2)
N(2)	32(2)	29(2)	34(2)	0(1)	9(1)	0(1)
C(2)	31(2)	30(2)	44(2)	2(2)	12(2)	4(2)
N(3)	29(2)	30(2)	29(2)	1(1)	5(1)	-3(1)
C(3)	32(2)	30(2)	46(2)	2(2)	10(2)	8(2)
C(4)	39(2)	51(2)	48(2)	2(2)	10(2)	5(2)
C(5)	51(3)	84(3)	43(2)	1(2)	3(2)	16(2)
C(6)	33(2)	82(3)	69(3)	-21(3)	6(2)	4(2)
C(7)	40(2)	75(3)	88(4)	1(3)	12(2)	-10(2)
C(8)	42(2)	61(3)	67(3)	15(2)	12(2)	-1(2)
C(9)	39(3)	166(6)	90(4)	-44(4)	-6(3)	2(3)
C(11)	36(2)	35(2)	26(2)	-3(2)	5(2)	4(2)
C(12)	39(2)	36(2)	42(2)	-6(2)	6(2)	-2(2)
C(13)	49(2)	36(2)	45(2)	-2(2)	8(2)	7(2)
C(14)	48(2)	45(2)	45(2)	-4(2)	8(2)	16(2)
C(15)	35(2)	47(2)	44(2)	-7(2)	6(2)	5(2)
C(16)	39(2)	33(2)	37(2)	-3(2)	7(2)	3(2)
C(17)	41(2)	47(2)	29(2)	-4(2)	-2(2)	7(2)
C(18)	48(2)	57(3)	37(2)	8(2)	-2(2)	10(2)
C(19)	41(2)	59(3)	51(3)	-5(2)	-9(2)	1(2)
C(21)	36(2)	25(2)	39(2)	0(2)	13(2)	-5(2)
C(22)	36(2)	35(2)	41(2)	-2(2)	9(2)	-5(2)
C(23)	54(2)	41(2)	45(2)	-6(2)	21(2)	-7(2)
C(24)	42(2)	50(3)	66(3)	-8(2)	28(2)	-9(2)
C(25)	38(2)	41(2)	61(3)	-6(2)	7(2)	-9(2)
C(26)	41(2)	35(2)	42(2)	3(2)	5(2)	-7(2)
C(27)	55(2)	29(2)	59(3)	0(2)	23(2)	3(2)
C(28)	85(3)	39(2)	96(4)	-11(2)	55(3)	-5(2)
C(29)	91(4)	44(3)	214(8)	-53(4)	89(5)	-25(3)
C(31)	35(2)	32(2)	27(2)	7(2)	2(2)	-7(2)
C(32)	43(2)	32(2)	48(2)	6(2)	5(2)	-1(2)
C(33)	59(3)	49(2)	34(2)	2(2)	7(2)	-21(2)
C(34)	46(2)	57(3)	40(2)	6(2)	-6(2)	-21(2)
C(35)	36(2)	51(2)	44(2)	13(2)	-3(2)	-7(2)
C(36)	34(2)	41(2)	41(2)	5(2)	5(2)	-3(2)
C(37)	35(2)	27(2)	36(2)	7(2)	9(2)	-4(2)

C(38)	33(2)	31(2)	51(2)	2(2)	8(2)	-5(2)
C(39)	55(2)	45(2)	43(2)	10(2)	18(2)	-6(2)
C(110)	73(3)	76(3)	32(2)	-5(2)	3(2)	21(3)
C(131)	66(3)	37(3)	112(4)	-3(3)	20(3)	6(2)
C(151)	39(2)	67(3)	96(4)	-8(3)	9(2)	8(2)
C(210)	174(7)	76(4)	86(4)	15(3)	12(4)	75(4)
C(231)	79(3)	82(3)	52(3)	-9(2)	35(2)	-17(3)
C(251)	40(2)	91(4)	88(4)	4(3)	0(2)	-9(2)
C(310)	43(2)	41(2)	34(2)	3(2)	8(2)	-7(2)
C(331)	80(3)	78(3)	53(3)	-27(3)	15(2)	-34(3)
C(351)	36(2)	76(3)	84(4)	5(3)	-5(2)	-3(2)

Table 7: Structural parameters for 1-OC(CH<sub>2</sub>)Ar<sup>2</sup> and 1-OC(CH<sub>2</sub>)NPhMe (crystallographically determined) and 2-OCHCH<sub>2</sub> and 2-OC(NH<sub>2</sub>)CH<sub>2</sub> (calculated).

	1-OC(CH <sub>2</sub> )Ar <sup>2</sup>	2-OCHCH <sub>2</sub>	1-OC(CH <sub>2</sub> )NPhMe	2-OC(NH <sub>2</sub> )CH <sub>2</sub>
Ti-O	1.826(2)	1.819	1.847(3)	1.828
C-O	1.352(4)	1.353	1.340(4)	1.360
Ti-N (avg.)	1.922	1.920	1.933	1.917
C=C (enol)	1.311(5)	1.337	1.309(5)	1.347
N-C (enol)			1.417(5)	1.396
O-Ti-N (avg.)	109	109	110	110
Ti-O-C	151.2(2)	151	159.0(2)	141
C(411)-C(41)-N(4)-C(48)			56	-22
C(411)-C(41)-N(4)-C(42)			-112	-50




Figure 5: Frontier molecular orbitals of 2-OCHCH<sub>2</sub> (top) and 2-OC(NH<sub>2</sub>)CH<sub>2</sub> (bottom).



1 DFT calculation for 2-OC(NH<sub>2</sub>)CH<sub>2</sub>

## Parallel Execution: Process Information

=====

Actual Number of Tasks running: 1  
The Master (kid 0) runs on host localhost  
(INPUT FILE)

title normal titanium enolate

atoms Z-matrix

1. c 0 0 0 0 0 0  
2. c 1 0 0 rCC 0 0  
3. o 2 1 0 rCO dd 0  
4. ti 3 2 1 rTiO aa 0  
5. n 4 3 2 rTiN bb 180  
6. n 4 3 2 rTiN bb 300  
7. n 4 3 2 rTiN bb 60  
8. h 1 2 3 rCH 120 180  
9. h 1 2 3 rCH 120 0  
10. n 2 1 9 rCN ee 180  
11. h 5 4 3 rNH cc 0  
12. h 5 4 3 rNH cc 180  
13. h 6 4 3 rNH cc 0  
14. h 6 4 3 rNH cc 180  
15. h 7 4 3 rNH cc 0  
16. h 7 4 3 rNH cc 180  
17 h 10 2 1 a ff fff  
18 h 10 2 1 a gg ggg  
end

geovar

rCC 1.31  
rCO 1.34  
rTiO 1.847  
rNH 1.02  
rTiN 1.93  
rCH 1.01  
rCN 1.41  
aa 159  
bb 109  
cc 120  
dd 122

Agapie et al.

Radical Scission of Symmetrical 1,4-Dicarbonyl Compounds

27

```
ee 120
a 1
ff 120
gg 120
fff 56
ggg -112
end
```

Symmetry=NOSYM

fragments

Ti t21.Ti

O t21.O

C t21.C

H t21.H

N t21.N

end

xc

LDA VWN

GRADIENTS BECKE PERDEW

end

relativistic zora

charge 0

corepotentials TAPE12 &

Ti 1

O 2

C 3

H 4

N 5

end

scf

ITERATIONS 99

MIXING 0.05

end

geometry

optim internal selected

end

end input

```
*****  
*  
* -----  
* Amsterdam Density Functional (ADF)          2000.01   17 May, 2000  
* -----  
*  
*  
*  
*  
*  
*  
* =====  
* |      A D F      |  
* |                  |  
* |                  |  
* =====  
*  
*  
* Online information and documentation: http://www.scm.com  
* E-mail: support@scm.com info@scm.com  
*  
* Scientific publications using ADF results must be properly referenced  
* See the User Manuals (or the web site) for recommended citations  
*  
***** pentium_linux *****
```

DF 2000.01 RunTime: Dec08-2000 20:26:57  
normal titanium enolate

A T T A C H E D     F I L E S

Core Potentials:  
file : TAPE12

## MODEL PARAMETERS

DENSITY FUNCTIONAL POTENTIAL (scf)

Agapie et al.

Radical Scission of Symmetrical 1,4-Dicarbonyl Compounds

29

LDA: VWN  
Gradient Corrections: Becke88 Perdew86

SPIN (restricted / unrestr.)  
Molecule: Restricted  
Fragments: Restricted

OTHER ASPECTS  
Relativistic Corrections: scalar (ZORA,FrozenCore)  
Core Treatment: Frozen Orbital(s)

Electric Field: ---  
Magnetic Field: ---

## Fragment File(s)

-----

c:

file : t21.C  
jobid: ADF 2000.01 RunTime: Dec08-2000 20:26:37  
title: Carbon (III, 1s frozen)

o:

file : t21.O  
jobid: ADF 2000.01 RunTime: Dec08-2000 20:26:30  
title: Oxygen (III, 1s frozen)

ti:

file : t21.Ti  
jobid: ADF 2000.01 RunTime: Dec08-2000 20:26:26  
title: Titanium (IV, 2p frozen)

n:

file : t21.N  
jobid: ADF 2000.01 RunTime: Dec08-2000 20:26:48  
title: Nitrogen (III, 1s frozen)

h:

file : t21.H  
jobid: ADF 2000.01 RunTime: Dec08-2000 20:26:41  
title: Hydrogen (III)

## 2 DFT calculation for 2-OCHCH<sub>2</sub>

### Parallel Execution: Process Information

=====

Actual Number of Tasks running: 1  
The Master (kid 0) runs on host localhost  
(INPUT FILE)

title normal titanium enolate

atoms Z-matrix

1. c 0 0 0 0 0 0  
2. c 1 0 0 rCC 0 0  
3. o 2 1 0 rCO dd 0  
4. ti 3 2 1 rTiO aa 0  
5. n 4 3 2 rTiN bb 180  
6. n 4 3 2 rTiN bb 300  
7. n 4 3 2 rTiN bb 60  
8. h 1 2 3 rCH 120 180  
9. h 1 2 3 rCH 120 0  
10. h 2 1 9 rCH ee 180  
11. h 5 4 3 rNH cc 0  
12. h 5 4 3 rNH cc 180  
13. h 6 4 3 rNH cc 0  
14. h 6 4 3 rNH cc 180  
15. h 7 4 3 rNH cc 0  
16. h 7 4 3 rNH cc 180  
end

geovar

rCC 1.31  
rCO 1.35  
rTiO 1.82  
rNH 1.2  
rTiN 1.92  
rCH 1  
aa 151.2  
bb 109  
cc 120  
dd 122  
ee 120  
end

Symmetry=C(s)

Agapie et al.

Radical Scission of Symmetrical 1,4-Dicarbonyl Compounds

31

fragments

Ti t21.Ti

O t21.O

C t21.C

H t21.H

N t21.N

end

xc

LDA VWN

GRADIENTS BECKE PERDEW

end

relativistic zora

charge 0

corepotentials TAPE12 &

Ti 1

O 2

C 3

H 4

N 5

end

scf

ITERATIONS 99

MIXING 0.05

end

geometry

optim internal selected

end

end input

```
* ****
```

```
* -----
```

```
* Amsterdam Density Functional (ADF)          2000.01   17 May, 2000
```

```
* -----
```

```
* 
```

```
* 
```

```
* 
```

```
* =====
```

```
* |              |
```

```
* |      A D F    |
```

```
* |              |
```

```
* =====
```

```
* 
```

```
* Online information and documentation: http://www.scm.com
```

```
* E-mail: support@scm.com info@scm.com
```

```
* 
```

```
* Scientific publications using ADF results must be properly referenced
```

```
* See the User Manuals (or the web site) for recommended citations
```

```
* 
```

```
* ***** pentium linux *****
```

ADF 2000.01 RunTime: Dec08-2000 00:59:38  
normal titanium enolate

A T T A C H E D     F I L E S

Core Potentials:  
file : TAPE12

## MODEL PARAMETERS

DENSITY FUNCTIONAL POTENTIAL (scf)

LDA:

VWN

Gradient Corrections: Becke88 Perdew86

SPIN (restricted / unrestr.)

Molecule: Restricted

Fragments: Restricted

OTHER ASPECTS

Relativistic Corrections: scalar (ZORA,FrozenCore)

Core Treatment: Frozen Orbital(s)

Electric Field: ---

Magnetic Field: ---

Fragment File(s)

c:

file : t21.C

jobid: ADF 2000.01 RunTime: Dec08-2000 00:59:25

title: Carbon (III, 1s frozen)

o:

file : t21.O

jobid: ADF 2000.01 RunTime: Dec08-2000 00:59:22

title: Oxygen (III, 1s frozen)

ti:

file : t21.Ti

jobid: ADF 2000.01 RunTime: Dec08-2000 00:59:13

title: Titanium (IV, 2p frozen)

n:

file : t21.N

jobid: ADF 2000.01 RunTime: Dec08-2000 00:59:36

title: Nitrogen (III, 1s frozen)

h:

file : t21.H

jobid: ADF 2000.01 RunTime: Dec08-2000 00:59:28

title: Hydrogen (III)